



## OPEN ACCESS

## EDITED BY

Ilene Sue Ruhoy,  
Mount Sinai South Nassau, United States

## REVIEWED BY

Sarah Cohen Solomon,  
PRISM Spine and Joint, United States

## \*CORRESPONDENCE

Alison M. Slater

✉ z5475629@ad.unsw.edu.au

S. Jade Barclay

✉ jade.barclay@sydney.edu.au

<sup>†</sup>These authors have contributed equally to this work and share first authorship

<sup>‡</sup>Senior authorship

RECEIVED 02 July 2024

ACCEPTED 22 July 2024

PUBLISHED 07 August 2024

## CITATION

Slater AM, Barclay SJ, Granfar RMS and Pratt RL (2024) Fascia as a regulatory system in health and disease.

*Front. Neurol.* 15:1458385.

doi: 10.3389/fneur.2024.1458385

## COPYRIGHT

© 2024 Slater, Barclay, Granfar and Pratt. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Fascia as a regulatory system in health and disease

Alison M. Slater<sup>1\*†</sup>, S. Jade Barclay<sup>2,3\*†</sup>, Rouha M. S. Granfar<sup>4</sup> and Rebecca L. Pratt<sup>5‡</sup>

<sup>1</sup>School of Population Health, The University of New South Wales, Sydney, NSW, Australia,

<sup>2</sup>Neuromuscular Imaging Research Lab, The Kolling Institute, St Leonards, NSW, Australia,

<sup>3</sup>Hypermobility and Performance Lab, Faculty of Medicine and Health, The University of Sydney, Sydney, NSW, Australia, <sup>4</sup>School of Pharmacy and Medical Sciences, Griffith University, Southport, QLD, Australia, <sup>5</sup>Department of Foundational Medical Studies, William Beaumont School of Medicine,

Rochester, MI, United States

Neurology and connective tissue are intimately interdependent systems and are critical in regulating many of the body's systems. Unlocking their multifaceted relationship can transform clinical understanding of the mechanisms involved in multisystemic regulation and dysregulation. The fascial system is highly innervated and rich with blood vessels, lymphatics, and hormonal and neurotransmitter receptors. Given its ubiquity, fascia may serve as a "watchman," receiving and processing information on whole body health. This paper reviews what constitutes fascia, why it is clinically important, and its contiguous and interdependent relationship with the nervous system. Unquestionably, fascial integrity is paramount to human locomotion, interaction with our environment, bodily sense, and general physical and emotional wellbeing, so an understanding of the fascial dysregulation that defines a range of pathological states, including hypermobility syndromes, autonomic dysregulation, mast cell activation, and acquired connective tissue disorders is critical in ensuring recognition, research, and appropriate management of these conditions, to the satisfaction of the patient as well as the treating practitioner.

## KEYWORDS

fascia, connective tissue, collagen dysregulation, hypermobility, neuroregulation, mast cell, inflammation

## 1 Introduction

Despite being declared a global public health priority, chronic musculoskeletal pain and dysregulation conditions remain under-funded, under-researched, misdiagnosed and grossly misunderstood (1). Research and clinical paradigms that dismiss the importance of fascia in pain regulation can lead to undertreatment and inappropriate treatment for pathological states such as hypermobile Ehlers-Danlos Syndrome (hEDS), autonomic dysregulation, mast cell activation, and acquired connective tissue disorders. Continuing to ignore the fascial system perpetuates misdiagnoses and inappropriate treatments, creating unnecessary prescriptions of ineffective medication. Furthermore, application of incongruous allied health management protocols can cause masking of symptoms, and contributes to increases in physician burnout, preventable disabilities, and the global burden of chronic pain management in healthcare.

Teaching the form and function of fascia is notably absent from most professional health care education and the lack of awareness limits current treatments. Conventional healthcare has historically overlooked fascia, framing it not as a system that facilitates and modulates sensory input but as inert filler material (2). Understanding fascial anatomy and its multifaceted

role as a regulatory system can transform how clinicians approach managing health and treating pain and disease.

By understanding the dynamics of the fascial system as it relates to the other organ systems of the body (musculoskeletal, neurovascular, endocrine, and immune)—we can illuminate its role in the regulation of pain, locomotion, inflammation, and autonomic conditions. People experiencing these conditions often have complex needs and difficult healthcare encounters, which can lead to clinical traumatization and mutual distrust between patients and clinicians (3). Continued research can help clinicians understand and improve care for “problem patients” by revealing an anatomical reason for their pain that travels and manifests in different ways on different days. This paper reviews the emerging body of evidence on fascia as a regulator of the body’s sensory input, and demonstrates how fascia-informed care can potentially improve patient outcomes and clinician experiences, ultimately preventing poor care across healthcare and medical specialties, particularly in the areas of neurovascular, hormonal, endocrine, musculoskeletal, inflammatory, and hypermobility-related conditions.

## 2 The regulatory role of fascia

### 2.1 Why fascia matters

Of the four major classes of tissues in the human body—endothelial, muscle, nerve, and connective tissue (fascia) (4)—fascia has historically been most overlooked and misunderstood in Western medicine (possibly due to ongoing nomenclature debates overshadowing clinically-relevant fascia studies (5)). For the purpose of this review, we will use the terms *fascia* and *connective tissue* interchangeably). Previously thought to be a passive material, the cellular membrane is now widely known to be a dynamic structure critical to cellular functionality. Similarly, fascia, once thought to be inert material, has now been found to host membrane-like receptors and perform active membrane-like functions for structures it connects and encapsulates (6). Current research indicates that fascia could be the moderating interface between many tissue types in the musculoskeletal, endocrine, and autonomic nervous systems. Fascia surrounds, supports and protects every nerve, muscle, blood vessel, and organ in the body, and is abundantly innervated (7, 8). Fascia is estimated to house over 250 million nerve endings (8), with sensory neurons outnumbering motor neurons 9:1 in some regions (9). Fascia houses 25% more nerve endings than skin, and 1,000% more than the collective innervation of muscle, so fascia could very well be considered our richest sensory organ (7, 8).

Connective (fascial) tissue manifests in many forms, from the loose connective tissue immediately under the skin, to the deep

connective tissue of the epimysium, perimysium and endomysium of muscle, to cartilage, tendon, ligament, to the protective wrapping around nerves and blood vessels, the periosteum and pericardium, and the visceral serous membrane. Traditionally discarded during anatomical dissection oriented toward muscles, organs, neurovascular bundles and bones, prominent clinicians and researchers note fascia’s emergence from obscurity as one of the most functionally diverse bodily components (2).

### 2.2 Functions of fascia

Fascia plays crucial roles in locomotion and regulation, and evolves in response to environmental stimuli and functional demand. Fascial integrity is paramount to movement, bodily sense, hormonal, autonomic and neurovascular regulation, and purposeful interaction with our environment. For movement and locomotion, fascia regulates posture (2), force transmission (10), strength generation (11), elastic recoil (12), proprioception (1), exteroception, and interoception (13). Fascia also regulates lymphatic efficacy (14), protection of delicate neural and vascular elements and organs (4), thermoregulation, inflammatory and immune responses (15), wound healing (16), hormonal production and secretion (adrenaline, estrogen, insulin, thyroid hormones, oxytocin) (17, 18), and venous return (19). Fascia plays a critical role in the transmission of neurotransmitters, namely serotonin, dopamine, GABA, and acetylcholine (17, 20). The regulation of peripheral resistance arteries by interfacing fascial tissue is essential for control of blood pressure, and for the increase in blood flow to the central nervous system and the heart under stress conditions (21). Alterations in the superficial fascia can also reciprocally lead to lymphedema/lipedema, which in turn may exacerbate suboptimal health of superficial fascia (22). Fascial morphology is initially determined by embryological and early childhood development (23), and undergoes perpetual adaptation across the lifespan in response to functional demand. Fascia is constantly evolving, both dependent on and modulating sustained postures, repetitive movement, quantity of motion, load (24), stress, strain, hydration (25), pH, temperature (23), neurotransmitters, and hormones (26).

### 2.3 Components of fascia

The fundamental components of fascial tissue are primarily specialized cells, collagen fibers, elastin fibers, and an extracellular matrix (ECM): (2).

- Cells provide the metabolic properties of the fascial tissue.
- Collagen and elastin fibers provide mechanical strength.
- ECM (also known as the ground substance) provides plasticity and elasticity.

Fascial cell types include fibroblasts, fasciocytes, adipocytes, macrophages and mast cells (27), undifferentiated mesenchyme cells, chondroblasts, chondrocytes, osteoblasts, and osteocytes (28). ECM contains hyaluronan (hyaluronic acid; HA), glycosaminoglycans (GAGs), water and ions (2). GAGs create an osmotic imbalance, which enables the ECM to attract up to 1,000 times its own weight in

Abbreviations: ADHD, Attention Deficit/Hyperactivity Disorder; ANS, Autonomic Nervous System; ECM, Extracellular Matrix; GAGs, Glycosaminoglycans; HA, Hyaluronic Acid; hEDS, Hypermobile Ehlers-Danlos Syndrome; HPA, Hypothalamus-Pituitary Axis; IMT, Intramuscular Connective Tissue; IV, Intravenous; IVIG, Intravenous Immunoglobulin; MCAD, Mast Cell Activation Disease; MCAS, Mast Cell Activation Syndrome; MRGPRX2, Mas-related G protein-coupled receptor-X2; NSAIDs, Non-steroidal anti-inflammatory drugs; POTS, Postural orthostatic tachycardia syndrome; SLE, Systemic lupus erythematosus; TGF-β1, Transforming growth factor-β1.

water (29). This hydrophilic quality of proteoglycans (glycoproteins that contain GAGs) is responsible for maintaining the volume of the extracellular matrix, which is constrained by the surrounding collagen fibers (30). The capacity for ECM to attract up to 1,000 times its weight in water is particularly important, because fascia is the “arena” in which localized acute inflammation and edema occurs (30).

Fascia evolves almost exclusively from the mesodermal layer during embryological development (23). Embryos first develop cerebrospinal fluid and fascia, then the remaining body systems and structures form within this “endless web” of fascia (23, 31). Fascia facilitates “a true continuity throughout our whole body” (2), and has been referred to as the “organ of form” (32), and “the architect of human movement” (22).

## 2.4 Functional demand specialization

Functional demand dictates the proportional distribution of each fascial component. Fascia develops perpetually across the lifespan in response to functional demand specific to each region of the body. Consider the specialization of three fascial components in response to specific functional demands:

- Type I collagen facilitates force transmission. For example, tendons and the thick aponeurotic sheets of the iliotibial band (33) and plantar fascia (34) facilitate force transmission (35, 36), so Type I collagen is the predominant fiber subtype.
- Fasciocytes facilitate sliding and gliding and secrete hyaluronan as a lubricant to facilitate movement. Where sliding and gliding is necessitated between muscle bellies, fasciocytes are distributed along the margins of the fascia that envelopes muscle (37), and hyaluronan is secreted to facilitate movement, particularly around the myofascial junction (38).
- Cartilage (also known as chondral tissue) is a highly specialized layer of dense ECM that allows almost frictionless motion in a joint. Cartilage is composed of hyaluronan, Type II collagen, and cartilage-specific proteoglycans interspersed sparsely with chondrocytes (39) enabling unique mechanical properties. Counterintuitively, “moderate mechanical loading” (40) on cartilage over time minimizes the turnover of tissue constituents, resulting in a protective effect rather than tissue degeneration (41).

## 2.5 Recognizing fascia dysregulation

Most clinical fascia literature focuses on temporary reversible impairment, assuming optimal homeostatic function in an otherwise healthy body. However, clinicians are often treating patients with (frequently undiagnosed) genetic variants altering the ECM fiber arrangement -- due to one or more changes affecting collagen proportions, quality, length, binding site affinities or distributions. In these cases, many of the “default” principles or “typical” assumptions do not apply, especially related to viscoelasticity, proprioception, nociception, elastic recoil and force transmission. When the usual ECM norms do not apply, the label “heterogenous” is often used to describe the diverse symptom presentations of individuals with conditions such as Hypermobility Ehlers-Danlos Syndrome (hEDS),

Marfan Syndrome, acquired connective tissue disease including systemic lupus erythematosus (SLE), polyarthralgia, systemic sclerosis, and Sjogren Syndrome to name a few.

Multimorbidity is the norm among individuals with collagen dysregulation, with an average of more than 10 diagnosed conditions in patients with hEDS (42, 43). Commonly reported conditions include general joint hypermobility (subluxations and dislocations of joints, tendons, and nerves, frequent soft tissue injury, and abnormal wound healing) (42–46), tethered cord and cervical instability (47), a range of oral and orofacial manifestations (48), inflammatory bowel disease (49), dysautonomia, irritable bowel syndrome, reflux, diverticulosis (50), and postural orthostatic tachycardia syndrome (POTS) (51); cardiovascular disease including mitral valve prolapse and aortic wall hyperelasticity (52); dermal hyperextensibility and atrophic scarring (44), easy bruising (53), asthma (54), lipedema/lymphedema, chronic pain (55, 56), anesthetic resistance (57), fibromyalgia, myalgic encephalomyelitis or chronic fatigue syndrome (50), central sensitization (58), abdominal hernias and pelvic organ prolapses (44). Mast cell activation syndrome (MCAS), hypermobility and dysautonomia have been identified as a triad to be aware of in clinical presentation (59), with higher odds of autism, ADHD (60, 61), panic disorder (62), anxiety diagnoses (63, 64), and neurodivergence (56).

## 3 Fascia, exercise and locomotion

### 3.1 Hyaluronan: lubricant or glue?

Hyaluronan (hyaluronic acid) is the primary constituent of the ECM (65). Hyaluronan is secreted as a lubricant to facilitate myofascial sliding and gliding of muscles and nerves. Hyaluronan has been found to contribute to cellular metabolism (66), morphogenesis (67), wound healing (68), and inflammation (69). In local anesthetic studies, hyaluronan has been noted to effect the efficacy and release rate of anesthesia (70), which may explain local anesthesia resistance reported by patients with collagen dysregulation conditions (57). Dysregulated hyaluronan is associated with cancer development, and can result in excessive tissue swelling, increased interstitial pressure and compression of neurovascular structures, causing pressure and pain (66, 71).

Hyaluronan dysregulation can occur either through under- or over-activity (67). Under-activity can lead to hyaluronan over-accumulation; over-activity can lead to hyaluronan over-production. Immobility precipitates pathological accumulation of hyaluronan, which super-aggregates in the tissues, impairing blood and lymphatic circulation, and reducing lymphatic efficacy (68). Excessive exercise may also overstimulate hyaluronan production, and subsequent super-aggregation causes it to act more like a glue than a lubricant (67). When hyaluronan acts like a glue, it can lead to symptoms of lipedema/lymphedema including fat tissue inflammation, painful adipose tissue, adipose tissue growth, and fibrosis (69, 72). This phenomenon of hyaluronan acting more like a glue than a lubricant has also been linked to delayed onset muscle soreness syndrome (DOMS) (73).

### 3.2 Muscles, posture and movement

Immediately beneath the dermal layers (of skin), the superficial fascia transects two adipose layers (the superficial adipose tissue and

deep adipose tissue). Superficial adipose tissue (SAT) regulates several aspects of whole-body physiology including insulin sensitivity, body temperature and immune responses (15). The adjacent superficial fascia plays a key role in the transmission of hormones and the protects invested vascular and neural plexuses (4). The superficial fascia is structured to support these circuitous structures, and facilitates the requisite flexibility to circumvent injury at the extremes of physiological range (74). The superficial fascia also includes the retinacula cutis, finger-like projections of collagen connecting the dermis above and hypodermis below, as well as the deep (muscular) fascia (75). The deep fascia comprises the epi-, peri- and endomysial layers of the muscular fascia. Collectively, these constitute the intramuscular connective tissue (IMT) (76), into which muscle fibers are wholly embedded (38, 77). With the seamless melding of the epi- and perimysial components, the IMT acts as a “scaffold” for muscle development and a carrier of the neural and vascular supply for muscle cells (76), in turn acting in series with the muscle itself and its associated mechanoreceptors. This allows forces generated within the locomotor system to be efficiently transmitted across joints (78), and so circumventing excessive articular strain.

Ruffini and Pacinian corpuscles are mechanoreceptors that appear in fascia in varying proportions (79). Ruffini corpuscles monitor persistent postural input, while Pacini receptors downregulate in response to continuous stimulus (80). “Myofascial expansions” (35)—wherein 30% of muscle fibers insert into fascia rather than a tendon (81)—facilitate force transmission (82) from both synergistic and antagonistic muscles into fascia rather than an enthesis. This accounts for 30% of the mechanical force (35, 83), helping to protect local neural and vascular tissue, as well as underlying joints (23), effectively allowing them to “float” (84). This calls into question some long-held anatomical paradigms around joint mechanics and load transfer in that muscle can no longer be considered the only element responsible for the organization of movement (35).

## 4 Fascia and regulation

### 4.1 Fascia and neuroregulation

The study of fascia demonstrates a specific distribution and precise localization of neural elements (8), closely connected with the central nervous system, and more so with the autonomic nervous system (ANS) (7). Local vasodilatation (80), and thermoregulation are primarily affected (85). In mast cell studies, fascia receptors have been found to respond to neurotransmitters more rapidly and at lower doses than neurons (86, 87). Treatments that target fascia regulation may have multiple uses. Current medical literature lacks fascia-focused research, and there is an identified need for studies designed specifically to evaluate the neuroregulation effects of treatments on fascia hyaluronan, and mast cells. Anxiety and depression have been associated with neuroinflammation, with mast cell stabilizers and treatments for inflammation yielding positive results (88). Limited clinical research has found ADHD medication to be effective in treating impulsivity and behavioral regulation in eating disorders (89), and has been reported in case studies to alleviate musculoskeletal and orofacial pain (90). The growing

understanding of the fundamental neural dynamics of fascial plasticity has catalyzed a paradigm shift in the manual therapy approach to treatment of fascial dysfunction, moving away from exclusive consideration of fascia’s mechanical properties, and calling into question the categorization of massage, tape, and compression as “passive therapies” (2). Chemical alterations of the extracellular matrix, and excessive mechanical stimulation are known to transform these receptors into nociceptors, which are appreciably more sensitive than the underlying musculature, sustaining longer-lasting hypersensitivity (79).

### 4.2 Inflammation, stress, anesthesia and pain

The dominance of sympathetic innervation has far-reaching consequences for fascial health and aspects of the hypothalamic–pituitary–adrenal (HPA) axis, given the abundance of autonomic neural input in this tissue (91, 92). Chronic stress and chronic pain are—jointly and individually—often accompanied by chronic inflammation (93, 94), with pro-inflammatory cytokines secreted by fibroblasts, myofibroblasts, adipocytes, mast cells, lymphocytes and vascular cells. As the fascia is the principal setting for inflammatory and immune system activity, fascial dysfunction can result in catastrophic cascades (79, 95, 96). Even under conditions of relatively low-grade chronic inflammation, cytokines denigrate the ECM, destabilizing connective tissue and stimulating fibrosis (87). Noradrenaline upregulates Transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1), which can cause fibroblasts to differentiate into myofibroblasts, which (in an environment of chronic stress and/or inflammation) then initiates fibrosis and contractures (97). Chronic adrenaline upregulation can similarly elicit contracture and inhibit effective wound healing (98), provoking structural adaptation. This in turn creates imbalance, pain and palpable tension (98–100). Stress hormones, like cortisol, and chronic inflammation can exert an adverse effect on muscle and bone quality (101, 102). The gut, with its enteric nervous system (little brain), is compartmentalized by fascia (103) and produces the same neurotransmitters as the brain. The vagus nerve purportedly acts as a bi-directional autonomic conduit between fascia and the central nervous system in countering sympathetic overload, as well as playing a key role in the gut-brain axis, tempering inflammation, facilitating intestinal homeostasis, satiety and energy regulation (104). The vagus nerve also connects the gut and the brainstem (36, 104), further strengthening the close relationship between fascial and the nervous systems.

Fascia appears to act as a mediator between the autonomic nervous system, emotional regulation and immune regulation (79). An emerging body of research has begun to explore the dynamic interplay between fascia and cancer, fascia and muscle function, fascia and neuroinflammation, dysautonomia, pain, hypermobility, and neurodiversity (56, 71, 76, 105), with some studies focused on the interaction of fascia, neurotransmitters, hormones, and the hypothalamic–pituitary–adrenal axis (HPA axis) (95, 98). Additionally, emerging research posits that fascia and hyaluronan play a key role in the perception of pain (1, 66, 106). Fascia is also richly endowed with endocannabinoid receptors, signifying its role as a source and modulator of pain, presenting possibilities for future pain management strategies and research (107).

While fascia can make inflammation and pain worse, facial regulation of inflammation can also provide an efficient therapeutic target to counteract chronic pain-inflammation-stress cascades. While MCAS is rarely the first diagnosis received (it is often the last of many), simple treatments targeting mast cell stabilization or inflammatory regulation in fascia like olopatadine (in Patanol eye drops or Patanase and Ryaltris nasal sprays), or sodium cromoglycate (also known as cromolyn sodium) have been reported to rapidly reduce these cascades, with topical or nasal application (86, 108–110).

Several drugs are associated with mast cell activation, including nonsteroidal anti-inflammatory drugs (NSAIDs), anesthetics, neuromuscular blocking agents and opiates/opioids, and contrasts used in radiology (87, 111). Approximately half of all perioperative drugs can trigger mast cell activation (111), and hyaluronan affects anesthesia efficacy and release rate (70), necessitating research into hyaluronan and mast cell receptors in connective tissue, and substances that bind with these receptors (or silence receptor-signaling) as therapeutic targets. One such receptor, Mas-related G protein-coupled receptor-X2 (MRGPRX2), binds with opioids and naltrexone, and has been associated with fibrosis, hypertrophic scarring, and Long COVID (87, 111). Low dose naltrexone (LDN) is thought to bind with these mast cell receptors, and research has begun to investigate its potential to reduce side effects of immunosuppressive therapies (86, 88, 110), enhance analgesia, reduce proinflammatory mediators, and efficacy as a treatment for MCAS, anxiety and depression, and neuropathic pain (110). After successful therapeutic responses in studies with back pain, plantar fasciitis and Long COVID (106, 112, 113), intravenous (IV) saline and IV immunoglobulin (IVIG) have been hypothesized to treat underlying fascia-immune dysregulation; which goes underrecognized and undertreated in a wide range of health conditions.

### 4.3 Mast cells and neuro-immuno-endocrine regulation

Mast cells (also known as mastocytes) are tissue-resident granulocytes of myeloid lineage that play a central role in adaptive and innate immune function, neurological and non-immunological processes, and pathologies far beyond allergy and mastocytosis. Mast cells—dubbed a “multi-functional master cell” (114)—are found in connective tissue, vascular tissue, adipose tissue, and lymphatic tissue around the body, and are highly concentrated and recruited to junctions where antigens could enter the body, including all mucosal openings, skin, blood, respiratory endothelium, and gastrointestinal tract (27, 114). In response to environmental stimuli, mast cell activation and degranulation modulates vasodilation, blood pressure, nociception, itch, fibrosis, tissue permeability, wound healing, inflammation and immune responses, and behavior (27, 114–116). Two distinct mast cell phenotypes have been identified, distinguished by granule content. Unlike mucosal mast cell granules that predominantly contain tryptase, mast cell granules of non-mucosal tissue mastocytes contain a wide range of specialized enzymes (114, 116, 117), and are rich pharmacological targets (118). When activated, mast cells release the content of the granules, which can have major local and systemic effects, particularly if mast cell derived mediators circulate via the vascular or lymphatic systems (86, 110).

Mast cells can also be activated in response to environmental changes, including temperature, pressure, and dermal vibration (119, 120). Mast cell activation contributes to swelling, itch, fasciitis, and predisposes an environment toward fibrosis via various mechanisms including fibroblast recruitment and proliferation, hyaluronan degradation, or excessive ECM deposition. With disruption or underlying dysregulation of connective tissue, a normal regulatory response may lead to a pathological outcome. When mast cells release enzymes and mediators (including histamine, cytokines, and hundreds more) (38, 116) in the neuromuscular-myofascial junctions, it may evoke an escalating cascade of multisystemic dysregulation and localized deterioration. It is worth noting that MCAS and mast cell activation disease (MCAD) are not interchangeable terms (121). The term MCAD includes several subtypes including systemic mastocytosis which is characterized by excessive quantities of mast cells, and MCAS which involves the inappropriate release/levels of mast cell chemicals. While the symptom presentation of these conditions is similar, the different pathomechanisms indicate different treatment plans. As mast cell activation increasingly features in cases of chronic pain-inflammation-stress cascades and anesthesia resistance within hypermobile populations and those with autonomic dysregulation (57, 59, 122), treatment plans that prioritize mast cell inhibition could prevent or reduce these cascades and increase the efficacy of multidisciplinary treatment protocols. This adds to the regulatory explanation for medication sensitivity and paradoxical responses to medications, and presents a strong case to target mast cells in clinical research and therapeutic intervention.

### 4.4 Hormonal regulation in muscular fascia

Research has discovered an abundance of estrogen and relaxin receptors in fascia, concentrated on the fibroblasts of muscular fascia (26). Hormone concentrations are known to fluctuate significantly throughout the menstrual cycle and pregnancy, declining with the onset of perimenopause (123). Hormones directly influence fascial stiffness and pain sensitization throughout the lifespan of people who menstruate, due to hormonal inhibition of fibrosis and inflammation resulting in ECM remodeling (25). This has major consequences in the consideration of athletes at any age, especially those with a hyperflexible fascial system, with or without the presence of joint hypermobility, and with or without the use of hormone therapies or oral contraceptives (124). People experiencing perimenopause and menopause may subsequently be more susceptible to fibromyalgia in the absence of premenopausal hormone concentrations (125).

## 5 Conclusion

### 5.1 Implications for practice and research

Dysregulation within the fascial system is considered scientifically complex and likely plays a pivotal role in the multimorbidity encountered by patients living with connective tissue diseases (diagnosed or not). Various stages of fascial health may lead to paradoxical or atypical responses to “normal” manual therapy, or pharmacotherapy; making walking or exercise painful or otherwise unmanageable by the patient. When therapy causes pain, fascial

impairment should be strongly considered and explored so treatment plans can better address the culprit of chronic pain. We call for clinical educators and researchers, and the next generation of healthcare practitioners to normalize the understanding of hyaluronan dysregulation and the fascial system and their ubiquitous role in whole body health, diagnosis and treatment planning, inflammation and pain management, and homeostasis.

## Author contributions

AS: Conceptualization, Writing – original draft, Writing – review & editing. JB: Conceptualization, Writing – original draft, Writing – review & editing. RG: Writing – review & editing. RP: Writing – original draft, Writing – review & editing.

## Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article. The author(s)

## References

- Langevin HM. Fascia mobility, proprioception, and myofascial pain. *Lifestyles*. (2021) 11:668. doi: 10.3390/life11070668
- Schleip R. Fascial plasticity - a new neurobiological explanation: part I. *J Bodyw Mov Ther*. (2003) 7:11–9. doi: 10.1016/S1360-8592(02)00067-0
- Halverson CME, Penwell HL, Francomano CA. Clinician-associated traumatization from difficult medical encounters: results from a qualitative interview study on the Ehlers-Danlos syndromes. *SSM - Qualitative Res Health*. (2023) 3:100237. doi: 10.1016/j.ssmqr.2023.100237
- Stecco C. Functional atlas of the human fascial system 1st Edition, Churchill Livingstone, Edinburgh: Elsevier Health Sciences (2015).
- Schleip R, Hedley G, Yucesoy CA. Fascial nomenclature: update on related consensus process. *Clin Anat*. (2019) 32:929–33. doi: 10.1002/ca.23423
- Fede C, Pirri C, Fan C, Petrelli L, Guidolin D, de Caro R, et al. A closer look at the cellular and molecular components of the deep/muscular fasciae. *Int J Mol Sci*. (2021) 22:1411. doi: 10.3390/ijms22031411
- Schleip R, and Stecco, C. 15 - Fascia as a Sensory Organ. In: R Schleip, J Wilke and A Baker, eds. *Fascia in Sport and Movement*. 2nd ed. (London, UK: Handspring Publishing) (2015) 169–180.
- Schleip R., Mechner, F., Zorn, A., and Klingler, W. (2014). The Bodywide Fascial Network as a Sensory Organ for Haptic Perception. *Journal of Motor Behavior*. 46, 191–193. doi: 10.1080/00222895.2014.880306
- Gesslbauer B, Hruby LA, Roche AD, Farina D, Blumer R, Aszmann OC. Axonal components of nerves innervating the human arm. *Ann Neurol*. (2017) 82:396–408. doi: 10.1002/ana.25018
- Tozzi P. A unifying neuro-fasciogenic model of somatic dysfunction – underlying mechanisms and treatment – part I. *J Bodyw Mov Ther*. (2015) 19:310–26. doi: 10.1016/j.jbmt.2015.01.001
- Schleip R, Muller DG. Training principles for fascial connective tissues: scientific foundation and suggested practical applications. *J Bodyw Mov Ther*. (2013) 17:103–15. doi: 10.1016/j.jbmt.2012.06.007
- Farris DJ, Raiteri BJ. Elastic ankle muscle-tendon interactions are adjusted to produce acceleration during walking in humans. *J Exp Biol*. (2017) 220:4252–60. doi: 10.1242/jeb.159749
- Schleip R, and Jäger, H. Interoception: A new correlate for intricate connections between fascial receptors, emotion, and self recognition. In: R Schleip, TW Findley, L Chaitow and PA Huijing, eds. *Fascia: The Tensional Network of the Human Body*. (London UK: Churchill Livingstone) (2012) 89–94. doi: 10.1016/B978-0-7020-3425-1.00047-7
- Albertin G, Astolfi L, Fede C, Simoni E, Contran M, Petrelli L, et al. Detection of lymphatic vessels in the superficial fascia of the abdomen. *Lifestyles*. (2023) 13:836. doi: 10.3390/life13030836
- Sakers A, De Siqueira MK, Seale P, Villanueva CJ. Adipose-tissue plasticity in health and disease. *Cell*. (2022) 185:419–46. doi: 10.1016/j.cell.2021.12.016
- Ye H, Rinkevich Y. Fascia layer—a novel target for the application of biomaterials in skin wound healing. *Int J Mol Sci*. (2023) 24:2936. doi: 10.3390/ijms24032936
- Schwartz A. Fascia and the vagus nerve. (2021). Available at: <https://drarielleschwartz.com/fascia-and-the-vagus-nerve-dr-arielle-schwartz/>
- Coelho M, Oliveira T, Fernandes R. Biochemistry of adipose tissue: an endocrine organ. *Archives of Med Sci: AMS*. (2013) 9:191–200. doi: 10.5114/aoms.2013.33181
- Bermudez K. Fasciotomy, chronic venous insufficiency, and the calf muscle pump. *JAMA*. (1999) 281:880.
- George T, De Jesus O. Physiology, Fascia. In: StatPearls. Treasure Island, FL: StatPearls Publishing (2024) - Available at: <https://www.ncbi.nlm.nih.gov/books/NBK568725/> (Accessed Mar 12, 2023).
- Storkebaum E, Carmeliet P. Paracrine control of vascular innervation in health and disease. *Acta Physiol*. (2011) 203:61–86. doi: 10.1111/j.1748-1716.2011.02333.x
- Kim Y, Park EY, Lee H. The effect of myofascial release in patients with breast cancer-related lymphedema: a cross-over randomized controlled trial. *Eur J Phys Rehabil Med*. (2023) 59:85–93. doi: 10.23736/S1973-9087.22.07698-5
- van der Wal JC. The architecture of the connective tissue in the musculoskeletal system - an often overlooked functional parameter as to proprioception in the locomotor apparatus. *Int J Therapeutic Massage and Bodywork*. (2009) 2:9–23. doi: 10.3822/ijmb.v2i4.62
- Zugel M, Maganaris CN, Wilke J. Fascial tissue research in sports medicine: from molecules to tissue adaptation, injury and diagnostics: consensus statement. *Br J Sports Med*. (2018) 52:1497. doi: 10.1136/bjsports-2018-099308
- Kodama Y, Masuda S, Ohmori T, Kanamaru A, Tanaka M, Sakaguchi T, et al. Response to mechanical properties and physiological challenges of fascia: diagnosis and rehabilitative therapeutic intervention for myofascial system disorders. *Bioengineering*. (2023) 10:474. doi: 10.3390/bioengineering10040474
- Fede C, Albertin G, Petrelli L, Sfriso MM, Biz C, de Caro R, et al. Hormone receptor expression in human fascial tissue. *European J Histochemistry: EJH*. (2016) 60:2710. doi: 10.4081/ejh.2016.2710
- Fede C, Petrelli L, Pirri C, Tiengo C, De Caro R, Stecco C. Detection of mast cells in human superficial fascia. *Int J Mol Sci*. (2023) 24:11599. doi: 10.3390/ijms241411599
- Świątek W, Kłodziński O, Brzeczek J, Kosiorowski I, Grzybowska N, Mozdziak PE. Components of the fascia – cells and extracellular matrix. *Medical J Cell Biol*. (2023) 11:13–9. doi: 10.2478/acb-2023-0002
- Wang ST, Neo BH, Betts RJ. Glycosaminoglycans: sweet as sugar targets for topical skin anti-aging. *Clin Cosmet Investig Dermatol*. (2021) 14:1227–46. doi: 10.2147/CCID.S328671

declare this research was partially funded through a grant from the Digital Health Cooperative Research Centre (DHCRC). The DHCRC is established and supported under the Australian Government's Cooperative Research Centres Program. Author JB holds a research scholarship funded through a DHCRC project grant.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

30. Langevin HM, Nedergaard M, Howe AK. Cellular control of connective tissue matrix tension. *J Cell Biochem.* (2013) 114:1714–9. doi: 10.1002/jcb.24521
31. Schultz, R. L., and Feitis, R. *The endless web: Fascial anatomy and physical reality.* Berkeley, CA: North Atlantic Books. (1996).
32. Varela FJ, Frenk S. The organ of form: towards a theory of biological shape. *J Soc Biol Struct.* (1987) 10:73–83. doi: 10.1016/0140-1750(87)90035-2
33. Hutchinson LA, Lichtwark GA, Willy RW, Kelly LA. The iliotibial band: a complex structure with versatile functions. *Sports Med.* (2022) 52:995–1008. doi: 10.1007/s40279-021-01634-3
34. Wilke J, Vleeming A, Wearing S. Overuse injury: the result of pathologically altered myofascial force transmission? *Exerc Sport Sci Rev.* (2019) 47:230–6. doi: 10.1249/JES.0000000000000205
35. Stecco A, Giordani F, Fede C, Pirri C, De Caro R, Stecco C. From muscle to the myofascial unit: current evidence and future perspectives. *Int J Mol Sci.* (2023) 24:4527. doi: 10.3390/ijms24054527
36. Swets, G. The power of fascia. National University Of Health Sciences: Chiropractic Medicine. [blog]. Available at: (2017). <https://www.nuhs.edu/the-power-of-fascia/>
37. Stecco C, Fede C, Macchi V, Porzionato A, Petrelli L, Biz C, et al. The fasciocytes: a new cell devoted to fascial gliding regulation. *Clin Anatomy (New York, NY).* (2018) 31:667–76. doi: 10.1002/ca.23072
38. Pirri C, Petrelli L, Guidolin D, Porzionato A, Fede C, Macchi V, et al. Myofascial junction: emerging insights into the connection between deep/muscular fascia and muscle. *Clin Anat.* (2024) 37:534–45. doi: 10.1002/ca.24148
39. Fox AJ, Bedi A, Rodeo SA. The basic science of articular cartilage. *Sports Health.* (2009) 1:461–8. doi: 10.1177/1941738109350438
40. Bader DL, Salter DM, Chowdhury TT. Biomechanical influence of cartilage homeostasis in health and disease. *Art.* (2011) 2011:979032:1–16. doi: 10.1155/2011/979032
41. Buckwalter JA, Mankin HJ, Grodzinsky AJ. Articular cartilage and osteoarthritis. *Instr Course Lect.* (2005) 54:465–80.
42. Petrucci T, Barclay SJ, Gensemer C, Morningstar J, Daylor V, Byerly K, et al. Phenotypic clusters and multimorbidity in hypermobile Ehlers-Danlos syndrome. *Mayo Clinic Proceed: Innov Quality & Outcomes.* (2024) 8:253–62. doi: 10.1016/j.mayocpiqo.2024.04.001
43. Demmler JC, Atkinson MD, Reinhold EJ, Choy E, Lyons RA, Brophy ST. Diagnosed prevalence of Ehlers-Danlos syndrome and hypermobility spectrum disorder in Wales, UK: a national electronic cohort study and case-control comparison. *BMJ Open.* (2019) 9:e031365. doi: 10.1136/bmjopen-2019-031365
44. Hakim, A. Hypermobile Ehlers-Danlos Syndrome. 2004 Oct 22 [Updated 2024 Feb 22]. In: Adam MP, Feldman J, Mirzaa GM, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle. (2004). Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1279/>.
45. Gensemer C, Burks R, Kautz S, Judge DP, Lavalley M, Norris RA. Hypermobile Ehlers-Danlos syndromes: complex phenotypes, challenging diagnoses, and poorly understood causes. *Dev Dyn.* (2021) 250:318–44. doi: 10.1002/dvdy.220
46. Dy CJ, Mackinnon SE. Ulnar neuropathy: evaluation and management. *Curr Rev Musculoskelet Med.* (2016) 9:178–84. doi: 10.1007/s12178-016-9327-x
47. Gensemer C, Daylor V, Nix J, Norris RA, Patel S. Co-occurrence of tethered cord syndrome and cervical spine instability in hypermobile Ehlers-Danlos syndrome. *Front Neurol.* (2024) 15:15. doi: 10.3389/fneur.2024.1441866
48. Lepperding U, Zschocke J, Kapferer-Seebacher I. Oral manifestations of Ehlers-Danlos syndromes. *Am J Med Genet C: Semin Med Genet.* (2021) 187:520–6. doi: 10.1002/ajmg.c.31941
49. Vounotrypidis P, Efremidou E, Zazos P, Pitiakoudis M, Maltezos E, Lyratzopoulos N, et al. Prevalence of joint hypermobility and patterns of articular manifestations in patients with inflammatory bowel disease. *Gastroenterol Res Pract.* (2009) 2009:924138:1–5. doi: 10.1155/2009/924138
50. Keer, R., and Grahame, R. (Eds.) (2003). *Hypermobility Syndrome. Recognition and Management for Physiotherapist.* London: Butterworth Heinemann. doi: 10.1016/B978-0-7506-5390-9.50006-X
51. Grigoriou E, Boris JR, Dormans JP. Postural orthostatic tachycardia syndrome (POTS): association with Ehlers-Danlos syndrome and Orthopaedic considerations. *Clin Orthop Relat Res.* (2015) 473:722–8. doi: 10.1007/s11999-014-3898-x
52. Mishra MB, Ryan P, Atkinson P. Extra-articular features of benign joint hypermobility syndrome. *Br J Rheumatol.* (1996) 35:861–6. doi: 10.1093/rheumatology/35.9.861
53. Edimo CO, Wajsborg JR, Wong S, Nahmias ZP, Riley BA. The dermatological aspects of hEDS in women. *Int J Women's Dermatol.* (2021) 7:285–9. doi: 10.1016/j.ijwd.2021.01.020
54. Morgan AW, Pearson SB, Davies S, Gooi HC, Bird HA. Asthma and airways collapse in two heritable disorders of connective tissue. *Ann Rheum Dis.* (2007) 66:1369–73. doi: 10.1136/ard.2006.062224
55. Ryan L, Beer H, Thomson E, Philcox E, Kelly C. Autistic traits correlate with chronic musculoskeletal pain: a self-selected population based survey. *OBM Neurobiol.* (2023) 7:1–21. doi: 10.21926/obm.neurobiol.2301155
56. Csesc JLL, Iodice V, Rae CL, Brooke A, Simmons R, Quadt L, et al. Joint hypermobility links Neurodivergence to Dysautonomia and pain. *Front Psychol.* (2022) 12:786916. doi: 10.3389/fpsy.2021.786916
57. Schubart JR, Schaefer E, Janicki P, Adhikary SD, Schilling A, Hakim AJ, et al. Resistance to local anesthesia in people with the Ehlers-Danlos syndromes presenting for dental surgery. *J Dental Anesthesia and Pain Med.* (2019) 19:261–70. doi: 10.17245/jdamp.2019.19.5.261
58. Schubert-Hjalmarsson E, Fasth A, Ickmans K, Mårdbrink EL, Söderpalm AC, Lundberg M. Central sensitization in adolescents with hypermobility spectrum disorder or hypermobile Ehlers-Danlos syndrome—a feasibility study. *Pilot and Feasibility Stud.* (2023) 9:97. doi: 10.1186/s40814-023-01320-3
59. Kucharik AH, Chang C. The relationship between hypermobile Ehlers-Danlos syndrome (hEDS), postural orthostatic tachycardia syndrome (POTS), and mast cell activation syndrome (MCAS). *Clin Rev Allergy Immunol.* (2020) 58:273–97. doi: 10.1007/s12016-019-08755-8
60. Baeza-Velasco C, Cohen D, Hamonet C, Vlamynck E, Diaz L, Cravero C, et al. Autism, joint hypermobility-related disorders and pain. *Front Psychol.* (2018) 9:656. doi: 10.3389/fpsy.2018.00656
61. Glans MR, Thelin N, Humble MB, Elwin M, Bejerot S. The relationship between generalised joint hypermobility and autism Spectrum disorder in adults: a large, cross-sectional, case control comparison. *Front Psychol.* (2022) 12:803334. doi: 10.3389/fpsy.2021.803334
62. Martin-Santos R, Bulbena A, Porta M, Gago J, Moilina L, Duro JC. Association between the joint hypermobility syndrome and panic disorder. *Am J Psychiatry.* (1998) 155:1578–83. doi: 10.1176/ajp.155.11.1578
63. Pailhez G, Bulbena A, Fullana MA, Castaño J. Anxiety disorders and joint hypermobility syndrome: the role of collagen tissue. *Gen Hosp Psychiatry.* (2009) 31:299. doi: 10.1016/j.genhosppsych.2008.08.009
64. Eccles JA, Owens AP, Mathias CJ, Umeda S, Critchley HD. Neurovisceral phenotypes in the expression of psychiatric symptoms. *Front Neurosci.* (2015) 9:4. doi: 10.3389/fnins.2015.00004
65. Pratt RL. Hyaluronan and the fascial frontier. *Int J Mol Sci.* (2021) 22:6845. doi: 10.3390/ijms22136845
66. Amir A, Kim S, Stecco A, Jankowski MP, Raghavan P. Hyaluronan homeostasis and its role in pain and muscle stiffness. *PM&R.* (2022) 14:1490–6. doi: 10.1002/pmrj.12771
67. Cowman MK, Schmidt TA, Raghavan P, Stecco A. Viscoelastic properties of Hyaluronan in physiological conditions. *F1000Research.* (2015) 4:622. doi: 10.12688/f1000research.6885.1
68. Stecco A, Cowman M, Pirri N, Raghavan P, Pirri C. Densification: Hyaluronan aggregation in different human organs. *Bioengineering.* (2022) 9:159. doi: 10.3390/bioengineering9040159
69. Duhon BH, Phan TT, Taylor SL, Crescenzi RL, Rutkowski JM. Current mechanistic understandings of lymphedema and lipedema: Tales of fluid, fat, and fibrosis. *Int J Mol Sci.* (2022) 23:6621. doi: 10.3390/ijms23126621
70. Shao P, Li H, Shi R, Li J, Wang Y. Understanding fascial anatomy and interfascial communication: implications in regional anesthesia. *J Anesth.* (2022) 36:554–63. doi: 10.1007/s00540-022-03082-3
71. Langevin HM, Keely P, Mao J, Hodge LM, Schleip R, Deng G, et al. Connecting (T)issues: how research in fascia biology can impact integrative oncology. *Cancer Res.* (2016) 76:6159–62. doi: 10.1158/0008-5472.CAN-16-0753
72. Tavianatou AG, Caon I, Franchi M, Piperigkou Z, Galessio D, Karamanos NK. Hyaluronan: molecular size-dependent signaling and biological functions in inflammation and cancer. *FEBS J.* (2019) 286:2883–908. doi: 10.1111/febs.14777
73. Wilke J, Behringer M. Is “delayed onset muscle soreness” a false friend? The potential implication of the fascial connective tissue in post-exercise discomfort. *Int J Mol Sci.* (2021) 22:9482. doi: 10.3390/ijms22179482
74. Guimberteau JC, Armstrong C. *Architecture of human living fascia: the extracellular matrix and cells revealed through endoscopy.* London, UK: Handspring Publishing (2015).
75. Tsukahara K, Tamatsu Y, Sugawara Y, Shimada K. Relationship between the depth of facial wrinkles and the density of the retinacula cutis. *Arch Dermatol.* (2012) 148:39–46. doi: 10.1001/archdermatol.2011.727
76. Purslow PP. The structure and role of intramuscular connective tissue in muscle function. *Front Physiol.* (2020) 11:495. doi: 10.3389/fphys.2020.00495
77. Trotter JA, Purslow PP. Functional morphology of the endomysium in series fibered muscles. *J Morphol.* (1992) 212:109–22. doi: 10.1002/jmor.1052120203
78. Purslow PP, Trotter JA. The morphology and mechanical properties of endomysium in series-fibered muscles: variations with muscle length. *J Muscle Res Cell Motil.* (1994) 15:299–308. doi: 10.1007/BF00123482

79. Schleip, R., and Bartsch, K. (2023). Fascia as a sensory and emotional organ: Fascia as a sensory organ. *Osteopathic medicine*. 24, 4–10. doi: 10.1016/S1615-9071(23)00009-6
80. Schleip R. Fascial plasticity – a new neurobiological explanation part 2. *J Bodyw Mov Ther.* (2003) 7:104–16. doi: 10.1016/S1360-8592(02)00076-1
81. Patel TJ, Lieber RL. Force transmission in skeletal muscle: from actomyosin to external tendons. *Exerc Sport Sci Rev.* (1997) 25:321–63. doi: 10.1249/00003677-199700250-00014
82. Stecco L. Atlas of physiology of the muscular fascia Padula, Italy: Piccin-Nuova Libreria (2016).
83. Huijting PA, van de Langenberg RW, Meesters JJ, Baan GC. Extramuscular myofascial force transmission also occurs between synergistic muscles and antagonistic muscles. *J Electromyogr Kinesiol.* (2007) 17:680–9. doi: 10.1016/j.jelekin.2007.02.005
84. Huijting PA. Muscle as a collagen fiber reinforced composite: a review of force transmission in muscle and whole limb. *Journal of Biomechanics.* (1999) 32:329–345. doi: 10.1016/S0021-9290(98)00186-9
85. Charkoudian N. Skin blood flow in adult human thermoregulation: how it works, when it does not, and why. *Mayo Clin Proc.* (2003) 78:603–12. doi: 10.4065/78.5.603
86. Molderings GJ, Haenisch B, Brettner S, Homann J, Menzen M, Dumoulin FL, et al. Pharmacological treatment options for mast cell activation disease. *Naunyn Schmiedeberg's Arch Pharmacol.* (2016) 389:671–94. doi: 10.1007/s00210-016-1247-1
87. Rama TA, Castells M. Triggers of anaphylaxis in Mastocytosis patients: evidence of the current drug-avoidance recommendation. *Curr Treat Options Allergy.* (2023) 10:442–57. doi: 10.1007/s40521-023-00349-2
88. McLaughlin PJ, Odom LB, Arnett PA, Orehek S, Thomas GA, Zagon IS. Low-dose naltrexone reduced anxiety in persons with multiple sclerosis during the COVID-19 pandemic. *Int Immunopharmacol.* (2022) 113:109438. doi: 10.1016/j.intimp.2022.109438
89. Keshen A, Bartel S, Frank GKW, Svedlund NE, Nunes A, Dixon L, et al. The potential role of stimulants in treating eating disorders. *Int J Eat Disord.* (2022) 55:318–31. doi: 10.1002/eat.23650
90. Zain E, Sugimoto A, Egawa J, Someya T. Case report: methylphenidate improved chronic pain in an adult patient with attention deficit hyperactivity disorder. *Front Psychol.* (2023) 14:1091399. doi: 10.3389/fpsy.2023.1091399
91. Fede C, Petrelli L, Pirri C, Neuherber W, Tiengo C, Biz C, et al. Innervation of human superficial fascia. *Front Neuroanat.* (2022) 16:981426–6. doi: 10.3389/fnana.2022.981426
92. Suarez-Rodriguez, V., Fede, C., Pirri, C., Petrelli, L., Loro-Ferrer, J. F., Rodriguez-Ruiz, D., et al. (2022). Fascial Innervation: A Systematic Review of the Literature. *Int. J. Mol. Sci.* 23:5674. doi: 10.3390/ijms23105674
93. Zhou, W. B. S., Meng, J., and Zhang, J. (2021). Does Low Grade Systemic Inflammation Have a Role in Chronic Pain? *Front Mol Neurosci.* 14:785214. doi: 10.3389/fmol.2021.785214
94. Ji RR, Xu ZZ, Gao YJ. Emerging targets in neuroinflammation-driven chronic pain. *Nat Rev Drug Discov.* (2014) 13:533–48. doi: 10.1038/nrd4334
95. Barsotti N, Chiera M, Lanaro D, Fioranelli M. Impact of stress, immunity, and signals from endocrine and nervous system on fascia. *Front Biosci-Elite.* (2020) 13:1–36. doi: 10.2741/870
96. Petrey AC, de la Motte CA. Hyaluronan, a crucial regulator of inflammation. *Front Immunol.* (2014) 5:101. doi: 10.3389/fimmu.2014.00101
97. Frangogiannis NG. Transforming growth factor- $\beta$  in tissue fibrosis. *J Exp Med.* (2020) 217:e20190103. doi: 10.1084/jem.20190103
98. Sivamani RK, Pullar CE, Manabat-Hidalgo CG, Rocke DM, Carlsen RC, Greenhalgh DG, et al. Stress-mediated increases in systemic and local epinephrine impair skin wound healing: potential new indication for Beta blockers. *PLoS Med.* (2009) 6:e1000012. doi: 10.1371/journal.pmed.1000012
99. Tozzi P. Does fascia hold memories? *J Bodyw Mov Ther.* (2014) 18:259–65. doi: 10.1016/j.jbmt.2013.11.010
100. Landén NX, Li D, Ståhle M. Transition from inflammation to proliferation: a critical step during wound healing. *Cell Mol Life Sci.* (2016) 73:3861–85. doi: 10.1007/s00018-016-2268-0
101. Molino A, Aversa Z, Muscaritoli M. Cortisol and the muscle-bone axis. *Osteoporos Int.* (2014) 25:2331–2. doi: 10.1007/s00198-014-2784-4
102. Hardy RS, Zhou H, Seibel MJ, Cooper MS. Glucocorticoids and bone: consequences of endogenous and exogenous excess and replacement therapy. *Endocr Rev.* (2018) 39:519–48. doi: 10.1210/er.2018-00097
103. Carabotti M, Scirocco A, Maselli MA, Severi C. The gut-brain axis: interactions between enteric microbiota, central and enteric nervous systems. *Annals Gastroenterol: Q Pub Hellenic Society of Gastroenterol.* (2015) 28:203–9.
104. Breit S, Kupferberg A, Rogler G, Hasler G. Vagus nerve as modulator of the brain-gut Axis in psychiatric and inflammatory disorders. *Front Psychol.* (2018) 9:44. doi: 10.3389/fpsy.2018.00044
105. Kerekes N, Sánchez-Pérez AM, Landry M. Neuroinflammation as a possible link between attention-deficit/hyperactivity disorder (ADHD) and pain. *Med Hypotheses.* (2021) 157:110717. doi: 10.1016/j.mehy.2021.110717
106. Schilder A, Hoheisel U, Magerl W, Benrath J, Klein T, Treede RD. Sensory findings after stimulation of the thoracolumbar fascia with hypertonic saline suggest its contribution to low back pain. *Pain.* (2014) 155:222–31. doi: 10.1016/j.pain.2013.09.025
107. Fede C, Pirri C, Petrelli L, Guidolin D, Fan C, de Caro R, et al. Sensitivity of the fasciae to the endocannabinoid system: production of Hyaluronan-rich vesicles and potential peripheral effects of cannabinoids in fascial tissue. *Int J Mol Sci.* (2020) 21:2936. doi: 10.3390/ijms21082936
108. Afrin LB, Butterfield JH, Raithel M, Molderings GJ. Often seen, rarely recognized: mast cell activation disease – a guide to diagnosis and therapeutic options. *Ann Med.* (2016) 48:190–201. doi: 10.3109/07853890.2016.1161231
109. Smith PK, Collins J. Olopatadine 0.6% nasal spray protects from vasomotor challenge in patients with severe vasomotor rhinitis. *American J Rhinol Allergy.* (2011) 25:e149–52. doi: 10.2500/ajra.2011.25.3620
110. Molderings GJ, Afrin LB. A survey of the currently known mast cell mediators with potential relevance for therapy of mast cell-induced symptoms. *Naunyn Schmiedeberg's Arch Pharmacol.* (2023) 396:2881–91. doi: 10.1007/s00210-023-02545-y
111. Navinés-Ferrer A, Serrano-Candelas E, Lafuente A, Muñoz-Cano R, Martín M, Gastaminza G. MRGPRX2-mediated mast cell response to drugs used in perioperative procedures and anaesthesia. *Sci Rep.* (2018) 8:11628. doi: 10.1038/s41598-018-29965-8
112. Peña-Martínez VM, Acosta-Olivo C, Tamez-Mata Y, Simental-Mendia LE, Blázquez-Saldaña J, Vilchez-Cavazos F, et al. Normal saline injection produces a therapeutic effect in patients with plantar fasciitis: a systematic review and meta-analysis of randomized controlled trials. *Foot Ankle Surg.* (2022) 28:1129–38. doi: 10.1016/j.fas.2022.04.005
113. Thompson JS, Thornton AC, Ainger T, Garvy BA. Long-term high-dose immunoglobulin successfully treats Long COVID patients with pulmonary, neurologic, and cardiologic symptoms. *Front Immunol.* (2023) 13:1033651. doi: 10.3389/fimmu.2022.1033651
114. Krystal-Whittemore M, Dileepan KN, Wood JG. Mast cell: a multi-functional master cell. *Front Immunol.* (2016) 6:620. doi: 10.3389/fimmu.2015.00620
115. da Silva EZM, Jamur MC, Oliver C. Mast cell function. *J Histochem Cytochem.* (2014) 62:698–738. doi: 10.1369/0022155414545334
116. Moon TC, Befus AD, Kulka M. Mast cell mediators: their differential release and the secretory pathways involved. *Front Immunol.* (2014) 5:569. doi: 10.3389/fimmu.2014.00569
117. Caughey GH. Mast cell tryptases and chymases in inflammation and host defense. *Immunol Rev.* (2007) 217:141–54. doi: 10.1111/j.1600-065X.2007.00509.x
118. Singh J, Shah R, Singh D. Targeting mast cells: uncovering prolific therapeutic role in myriad diseases. *Int Immunopharmacol.* (2016) 40:362–84. doi: 10.1016/j.intimp.2016.09.019
119. Li H, Guo Z, Xiangdong Q. Role of mechanical stimulus in mast cell activation. *Digital Med.* (2024) 10:e23. doi: 10.1097/DM-2023-00014
120. Boyden SE, Desai A, Cruse G, Young ML, Bolan HC, Scott LM, et al. Vibratory Urticaria associated with a missense variant in ADGRE2. *N Engl J Med.* (2016) 374:656–63. doi: 10.1056/NEJMoa1500611
121. Molderings GJ, Brettner S, Homann J, Afrin LB. Mast cell activation disease: a concise practical guide for diagnostic workup and therapeutic options. *J Hematol Oncol.* (2011) 4:10. doi: 10.1186/1756-8722-4-10
122. Blitshteyn S. Dysautonomia, hypermobility Spectrum disorders and mast cell activation syndrome as migraine comorbidities. *Curr Neurol Neurosci Rep.* (2023) 23:769–76. doi: 10.1007/s11910-023-01307-w
123. Sims ST, Yeager S. Next Level. Your guide to kicking ass, feeling great, and crushing goals through menopause and beyond. New York: Rodale Books (2022).
124. Vita M, Sedlackova Z, Herman M, Furst T, Smekal D, Cech Z. Influence of female hormones on fascia elasticity: an elastography study. *Clin Anat.* (2019) 32:941–7. doi: 10.1002/ca.23428
125. Ruschak I, Montesó-Curto P, Rosselló L, Aguilar Martín C, Sánchez-Montesó L, Toussaint L. Fibromyalgia syndrome pain in men and women: a scoping review. *Health.* (2023) 11:223. doi: 10.3390/healthcare11020223